



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GENERAL INDEX TO VOLUME V

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italic*; and previously published scientific names and all other matter, in ordinary type.

A

- abnormis* (*Hymenochaete*), 367
acerina (*Thelephora*), 196
acerinum (*Corticium*), 196
acerinum (*Stereum*), 196
acerinum var. *nivosum* (*Stereum*), 193
acerinus (*Aleurodiscus*), 196
agglutinans (*Hymenochaete*), 344
Aleurodiscus, 177; *acerinus*, 196, *amorphus*, 177, 180; *apiculatus*, 186; *botryosus*, 198; *candidus*, 188; *cremeus*, 199; *Farlowii*, 182; *Mancinellus*, 190; *nivosus*, 193; *Oakesii*, 183; *penicillatus*, 201; *seriatus*, 192; *strumosus*, 190; *tenuis*, 200; *Weirii*, 203
Alga, a wood-penetrating, 211
Algological notes. III. A wood-penetrating alga, *Gomontia lignicola*, n. sp., 211
ambiens (*Hymenochaete*), 344
amorpha (*Cyphella*), 180
amorpha (*Peziza*), 180
amorpha (*Thelephora*), 180
amorphum (*Corticium*), 180
amorphus (*Aleurodiscus*), 177, 180
anomala (*Hymenochaete*), 358
apiculatus (*Aleurodiscus*), 186
arida (*Hymenochaete*), 340
arida (*Hymenochaetella*), 340
aspera (*Hymenochaete*), 311, 370
asperata (*Hymenochaete*), 351
Auricularia ferruginea, 332; *tabacina*, 325
avellana (*Hymenochaete*), 325
avellana (*Thelephora*), 325
avellanum (*Stereum*), 325

B

- badio-ferruginea* (*Hymenochaete*), 330, 370
badio-ferrugineum (*Stereum*), 330
balsamicola (*Nodularia*), 180
Berkeleyana (*Hymenochaete*), 313, 370
Berkeleyanum (*Stereum*), 313
Biscutella, 147

- Bonns, W. W.: Etherization of tissues and its effect on enzyme activity, 225; Duggar, B. M., and. The effect of Bordeaux mixture on the rate of transpiration, 153
Bordeaux mixture, The effect of, on the rate of transpiration, 153
borealis (*Hymenochaete*), 317, 370
botryosum (*Corticium*), 128
botryosus (*Aleurodiscus*), 198
Burt, E. A. *Corticiums* causing Pellicularia disease of the coffee plant, hypochnose of pomaceous fruits, and Rhizoctonia disease, 119; The *Thelephoraceae* of North America, IX, 177; X, 301

C

- Cacao (*Hymenochaete*), 310
Cacao (*Stereum*), 310
candida (*Thelephora*), 188
candidissima (*Thelephora*), 188
candidum (*Stereum*), 188
candidus (*Aleurodiscus*), 188
Castor bean leaves, transpiration of, 171
cervina (*Hymenochaete*), 363
Chamonixia caespitosa, 140
Chemical reactions, effect of anaesthetics on, 251
Choeromyces meandriformis, 29
Cineraria heterophylla, 72; *integrifolia* β *minor*, 70, 74; *vulneraria*, 91
cinnamomea (*Hymenochaete*), 345
cinnamomea (*Thelephora*), 345
cinnamomeum (*Corticium*), 345
Correlation of the strength and durability of southern pine, 109
corrugata (*Hymenochaete*), 358
corrugata (*Thelephora*), 358
corrugatum (*Corticium*), 358
Corticium acerinum, 196; *amorphum*, 180; *botryosum*, 128; *cinnamomeum*, 345; *corrugatum*, 358; *epichlorum*, 351; *koleroga*, 123; *Oakesii*, 183; *ochroleucum*, 125; *simulans*, 341;

(373)

simulans, 340; *Solani*, 128; **Stevensii**, 125; *vagum*, 127, 128; *vagum* var. *Solani*, 128
 Corticiums causing Pellicularia disease of the coffee plant, hypochnose of pomaceous fruits, and Rhizoctonia disease, 119
corticolor (Hymenochaete), 339, 370
crassa (Hymenochaete), 367
cremeus (Aleurodiscus), 199
 Cruciferae, Notes on certain, 143
cubensis (Hymenochaete), 337, 372
Curtisii (Hymenochaete), 320, 372
Curtisii (Stereum), 320
Cyperus, transpiration of, 168
Cyphella amorpha, 180

D

damaecornis (Hymenochaete), 306, 370
damaecornis (Thelephora), 306
damicornae (Stereum), 306
digitata (Hymenochaete), 347
Dithyrea, 147; *californica*, 150, var. *maritima*, 150; *clinata*, 150; *Griffithsii*, 148; *membranacea*, 149; *Wislizenii*, 147, var. *Griffithsii*, 148, var. *Palmeri*, 148
 Dodge, C. W., Zeller, S. M., and. Gautieria in North America, 133; Rhizopogon in North America, 1
Draba *Gilgiana*, 151; *gilgiana*, 150; **Standleyi**, 150
 Duggar, B. M., and Bonns, W. W. The effect of Bordeaux mixture on the rate of transpiration, 153
dura (Hymenochaete), 352
 Durability of pine, 109

E

Effect of Bordeaux mixture on the rate of transpiration, The, 153
elegantissima (Hymenochaete), 314
elegantissimum (Stereum), 314
 Enzyme activity, effect of etherization of tissues on, 225
epichlora (Hymenochaete), 351
epichlorum (Corticium), 351
episphaeria (Hymenochaete), 362
episphaeria (Thelephora), 362
Erysiphe scandens, 123
 Etherization of tissues and its effect on enzyme activity, 225

F

Farlowii (Aleurodiscus), 182
ferruginea (Auricularia), 332
ferruginea (Hymenochaete), 332
ferrugineum (Stereum), 332

fimbriata (Hymenochaete), 367
Foreliella, 212; *perforans*, 212
formosa (Hymenochaete), 308, 370
formosa (Hymenochaete), 307
frustulosa (Hymenochaete), 367
frustulosum (Septobasidium), 367
fuliginosa (Hymenochaete), 342, 365, 372
fuliginosa (Hymenochaete), 342
fuliginosa (Thelephora), 365
fuliginosum (Stereum), 365
fulva (Hymenochaete), 354
fulvella (Hymenochaete), 318
fusca (Hymenochaete), 366
fusca (Hymenochaetella), 366

G

Gautiera, 133
Gautieria *Drummondii*, 141; *graveolens*, 136, 142; *graveolens*, 136, var. *mexicana*, 136; *monticola*, 139, 142; *morchelliformis*, 134, 142; *morriellaeformis*, 135; *Othii*, 141; **plumbea**, 138, 142; *Trabuti*, 137, 142; *villosa*, 135
Gautieria in North America, 133
Gomontia *Aegagropilae*, 212; *arrhiza*, 212; *codiolifera*, 212; *Holdenii*, 212; *lignicola*, 211, 218, 220, 222, 224; *Manxiana*, 212; *perforans*, 212; *polyrhiza*, 212
 Greenman, J. M.: Monograph of the North and Central American species of the genus *Senecio*—Part II, 37; and Pfeiffer, Norma E. A new selaginella from Mexico, 205

H

Helvella nicotiana, 325; *rubiginosa*, 332
Hydnangium aurantium, 30
Hydnochaete setosa, 312
Hydnum resupinatum, 311
 Hymenochaete, 301; *abnormis*, 367; *agglutinans*, 344; *ambiens*, 344; **anomala**, 358; *arida*, 340; *aspera*, 311, 370; *asperata*, 351; *avellana*, 325; *radio-ferruginea*, 330, 370; *Berkkeleyana*, 313, 370; *borealis*, 317, 370; *Cacao*, 310; *cervina*, 363; *cinnamomea*, 345; *corrugata*, 358; *corticolor*, 339, 370; *crassa*, 367; **cubensis**, 337, 372; *Curtisii*, 320, 372; *damaecornis*, 306, 370; *digitata*, 347; *dura*, 352; *elegantissima*, 314; *epichlora*, 351; *episphaeria*, 362; *ferruginea*, 332; *fimbriata*, 367; *formosa*, 308, 370; *formosa*, 307; *frustulosa*, 367; *fuliginosa*, 342, 365, 372; *fuliginosa*, 342; **fulva**, 354;

- fulvella*, 318; *fusca*, 366; *imbricata*, 325; *imbricatula*, 325; *insularis*, 358; *Kunzei*, 323; *laeta*, 323; *laevigata*, 348; *laxa*, 340; *leonina*, 353; *luteo-badia*, 323, 372; **multisetae**, 357; *multispinulosa*, 367; *musicola*, 367; *opaca*, 364; *pallida*, 367; *paupercula*, 367; *pinnatifida*, 355, 372; *pulcherrima*, 318; *purpurea*, 368; **reflexa**, 336, 372; *reniformis*, 310; *rigidula*, 318; *rubiginosa*, 332, 372; *rudis*, 346; *rugispora*, 367; *Sallei*, 314, 372; *Schomburgkii*, 308; *scruposa*, 318; *setosa*, 311, 368; *simulans*, 341; *siparia*, 368; *spretia*, 348; *tabacina*, 325, 372; *tenuis*, 364; *tenuissima*, 315; *tenuissima*, 314; *tomentosa*, 368; *ungulata*, 338, 372; *unicolor*, 342
- Hymenochaetella arida*, 340; *fusca*, 366; *laxa*, 340; *rudis*, 346
- Hymenogaster Klotzschii*, 29; *monticolus*, 139; *Trabuti*, 137
- Hypochnopsis ochroleuca*, 125
- Hypochnose of pomaceous fruits, 125
- Hypochnus ochroleucus*, 125; *Solani*, 128
- Hysterangium Duriaeanum*, 11; *rubescens*, 18
- Hysteromyces*, 1; *graveolens*, 15

I

- imbricata* (*Hymenochaete*), 325
- imbricatula* (*Hymenochaete*), 325
- imbricatula* (*Thelephora*), 325
- insularis* (*Hymenochaete*), 358

K

- koleroga* (*Corticium*), 123
- koleroga* (*Pellicularia*), 123
- Kunzei* (*Hymenochaete*), 323
- Kunzei* (*Thelephora*), 323

L

- laeta* (*Hymenochaete*), 323
- laetum* (*Stereum*), 323
- laevigata* (*Hymenochaete*), 348
- laxa* (*Hymenochaete*), 340
- laxa* (*Hymenochaetella*), 340
- Lenzites saepiaria*, 111
- leonina* (*Hymenochaete*), 353
- Lesquerella*, 143; *lasiocarpa*, 143
- luteo-badia* (*Hymenochaete*), 323, 372
- luteo-badia* (*Thelephora*), 323
- luteo-badium* (*Stereum*), 323
- Lycoperdon aestivum*, 28

M

- Mancianum* (*Stereum*), 190
- Mancianus* (*Aleurodiscus*), 190
- Marguerites, transpiration of, 165
- Melanogaster Berkeleyanus*, 18
- Metabolism, effect of anaesthetics on, 243
- Moore, George T. Algological notes.
- III. A wood-penetrating alga, *Gomontia lignicola*, n. sp., 211
- multisetae** (*Hymenochaete*), 357
- multispinulosa* (*Hymenochaete*), 367
- musicola* (*Hymenochaete*), 367
- Mylitta roseola*, 16

N

- Narcosis, stimulation, inhibition, and the theory of, 253
- nicotiana* (*Helvella*), 325
- nivosus* (*Aleurodiscus*), 193
- Nodularia balsamicola*, 180
- Notes on certain Cruciferae, 143

O

- Oakesii* (*Aleurodiscus*), 183
- Oakesii* (*Corticium*), 183
- ochroleuca* (*Hypochnopsis*), 125
- ochroleucum* (*Corticium*), 125
- ochroleucus* (*Hypochnus*), 125
- opaca** (*Hymenochaete*), 364

P

- pallida* (*Hymenochaete*), 367
- paupercula* (*Hymenochaete*), 367
- Payson, E. B. Notes on certain Cruciferae, 143
- Pellicularia* disease of coffee plant, 119
- Pellicularia koleroga*, 123
- penicillatus** (*Aleurodiscus*), 201
- Permeability, effect of anaesthetics on, 240
- Peziza amorpha*, 180
- Pfeiffer, Norma E., Greenman, J. M. and. A new selaginella from Mexico, 205
- Photosynthesis, effect of anaesthetics on, 235
- Physaria*, 143; *acutifolia*, 145; *alpestris*, 147; *brassicoides*, 145; *didymocarpa*, 144, var. *australis*, 144, var. *lanata*, 145; *didymocarpa*, 145, var. *Newberryi*, 146; *floribunda*, 146; *Geyeri*, 146; *lanata*, 145; *Newberryi*, 146; *oregona*, 146; *Osterhoutii*, 146; *vitulifera*, 145
- Pine, southern, Correlation of the strength and durability of, 109
- pinnatifida** (*Hymenochaete*), 355, 372

Pinus echinata, 110, original samples of, 116; *palustris*, 109, 110, original samples of, 118
 Potatoes, transpiration of, 158
pulcherrima (*Hymenochaete*), 318
pulchrum (*Stereum*), 323
purpurea (*Hymenochaete*), 368

R

reflexa (*Hymenochaete*), 336, 372
reniforme (*Stereum*), 310
reniformis (*Hymenochaete*), 310
 Respiration, effect of anaesthetics on, 235
resupinatum (*Hydnum*), 311
 Rhizoctonia disease, 127
Rhizoctonia Solani, 129
Rhizopogon, 1; *aestivus*, 28; *albus*, 29; *albus*, 27, 29; *angustisepta*, 24, 36; *aurantius*, 30; *borealis*, 28; Briardi, 25; *diplophloeus*, 8, 34; *dubius*, 29; *graveolens*, 15, 32; *induratus*, 17, 36; *lapponicus*, 27; *Leonis*, 30; *luteolus*, 1, 10, 32; *luteolus*, 18, 27; *maculatus*, 4, 32, 34; *occidentalis*, 14, 32, 34; *pachyphloeus*, 9, 32, 34; *pannosus*, 6, 32, 34; *piceus*, 28; *provincialis*, 13, 32; *Rodwayi*, 22; *roseolus*, 16, 32, 36; *rubescens*, 18, var. *Vittadinii*, 20, 32; *rubescens*, 16; *rubrocorticeus*, 25, 36; *suavis*, 23; *violaceus*, 21; *virens*, 28; *virens*, 26; *virescens*, 26; *viridis*, 5; *Webbii*, 27
Rhizopogon in North America, 1
rigidula (*Hymenochaete*), 318
 Rotating table used in transpiration experiments, 154, 176
rubiginosa (*Helvella*), 332
rubiginosa (*Hymenochaete*), 332, 372
rubiginosa (*Thelephora*), 332
rubiginosum (*Stereum*), 332
rudis (*Hymenochaete*), 346
rudis (*Hymenochaetella*), 346
rugispora (*Hymenochaete*), 367

S

Sallei (*Hymenochaete*), 314, 372
scandens (*Erysiphe*), 123
Schomburgkii (*Hymenochaete*), 308
scruposa (*Hymenochaete*), 318
Selaginella, a new, from Mexico, 205
Selaginella Landii, 205, 208, 210; *rupestris*, 206; *Watsoni*, 206
 Senecio, Monograph of the American species of the genus—Part II, 37
 Senecio, 37; *Actinella*, 63, var. *mogollonicus*, 64; *Alabamensis*, 70; *alpicola*, 58; *altus*, 101; *antennariifolius*, 72; *appendiculatus*, 47; *arizonicus*, 40; *arizonicus*, 43; *atratus*, 97, 108,

var. *milleflorus*, 100; *aurellus*, 48; *aureus* var. *alpinus*, 58, var. *borealis*, 41, 56, 86, var. *obovatus*, 68, var. *werneriaefolius*, 60; *bellidifolius*, 91; *bernardinus*, 45, var. *sparsilobatus*, 46; *Blumeri*, 41; *candidissimus*, 90; *canovirens*, 55; *canus*, 73; *canus*, 78, 80, 84, 88, var. *acraeus*, 74, var. *celsus*, 74, var. *Purshianus*, 80, var. *pygmaea*, 63, var. *Rothrock*, 78; *cheiranthifolius*, 91; *cognatus*, 48; *collinus*, 80; *confertus*, 50; *convallium*, 67; *cynthioides*, 93; *Encelesia*, 41; *eurypterus*, 47; *exaltatus* var. *minor*, 98; *fastigiatus*, 94, var. *Macounii*, 96; *fastigiatus*, 93, var. *eroseo-dentatus*, 96; *Fendleri*, 49, var. *canovirens*, 55, var. *molestus*, 54; *Fendleri* var. *lanatus*, 50, var. *subintegra*, 93; *gerberaefolius*, 66; *Greenei*, 67; *Hallii*, 90, 104; *Harrowii*, 78; *Hartmanii*, 44; *Howellii*, 83, var. *eradiatus*, 87, var. *lithophilus*, 86; *integrifolius*, 73, β . *heterophylla*, 72; *ionophyllus* var. *bernardinus*, 45; *kernensis*, 88; *lanatifolius*, 50; *Laramiensis*, 80; *Leonardii*, 68; *leucoorinus*, 94; *ligulifolius*, 94; *loratifolius*, 93, 106; *lugens* var. *exaltatus*, 98, 100, var. *foliosus*, 98; *Macounii*, 96; *milleflorus*, 100; *mogollonicus*, 65; *molinarius*, 65; *Muirii*, 56; *mutabilis*, 48; *Nelsonii*, 49; *neo-mexicanus*, 41, 104, var. *Griffithsii*, 43; *neo-mexicanus*, 45, 47; *oreophilus*, 44; *oreopolus*, 88, forma *aphanactis*, 89; *oresbius*, 44; *pauciflorus*, 91; *pentodontus*, 58; *perennans*, 60; *petraeus*, 57; *petrocallis*, 58; *petrophilus*, 58; *Purshianus*, 79; *Purshianus*, 73, 84, var. *viridescens*, 73; *rosulatus*, 49; *rosulatus* var. *coryphocolus*, 50, mut. *primulinus*, 50; *Rothrockii*, 78; *salcinus*, 49; *saxosus*, 57; *scaposus*, 60; *sordidus*, 41; *sparsilobatus*, 46; *spatuliformis*, 94; *sphaerocephalus*, 101; *Thurberi*, 62; *tomentosus*, 69; *tomentosus* var. β , 72; *Toumeyii*, 41; *trichophalus*, 91; *turbinatus*, 58; *umbraclifer*, 97; *Vulneraria*, 91; *werneriaefolius*, 60, var. *incertus*, 62; *werneriaefolius*, 56; *willowensis*, 41; *Wrightii*, 93

Septobasidium frustulosum, 367
seriatum (*Stereum*), 192
seriatum (*Aleurodiscus*), 192
setosa (*Hydnochaete*), 312
setosa (*Hymenochaete*), 311, 368
setosa (*Thelephora*), 311
simulans (*Corticium*), 341
simulans (*Corticium*), 340

- simulans* (*Hymenochaete*), 341
siparia (*Hymenochaete*), 368
Solani (*Corticium*), 128
Solani (*Hypochnus*), 128
Solani (*Rhizoctonia*), 129
speciosa (*Thelephora*), 307
Splanchnomyces, 1; *Cauvinianus*, 11;
dubius, 29; *luteolus*, 11; *Rabenhorstii*, 11; *roseolus*, 16; *Webbiana*,
 27
spretia (*Hymenochaete*), 348
Stereum acerinum, 196, var. *nivosum*,
 193; *avellanum*, 325; *badio-ferru-*
gineum, 330; *Berkeleyanum*, 313;
Cacao, 310; *candidum*, 188; *Curtisii*,
 320; *damicorne*, 306; *elegantissimum*,
 314; *ferrugineum*, 332; *fuliginosum*,
 365; *laetum*, 323; *luteo-badium*, 323;
Mancianum, 190; *pulchrum*, 323;
reniforme, 310; *rubiginosum*, 332;
seriatum, 192; *strumosum*, 190;
tabacinum, 325; *tenuissimum*, 314;
tenuissimum, 314; *umbrinum*, 367,
 368; *vitellinum*, 190
Stevensii (*Corticium*), 125
 Strength and durability of yellow pine,
 109
strumosum (*Stereum*), 190
strumosus (*Aleurodiscus*), 190
Synthlipsis, 143; *Berlandieri*, 143;
Greggii, 143

T

- tabacina* (*Auricularia*), 325
tabacina (*Hymenochaete*), 325, 372
tabacina (*Thelephora*), 325
tabacinum (*Stereum*), 325
tenuis (*Aleurodiscus*), 200
tenuis (*Hymenochaete*), 364
tenuissima (*Hymenochaete*), 314
tenuissima (*Hymenochaete*), 315
tenuissimum (*Stereum*), 314
tenuissimum (*Stereum*), 314
Terfezia Leonis, 30

- Thelephora acerina*, 196; *amorpha*,
 180; *avellana*, 325; *candida*, 188;
candidissima, 188; *cinnamomea*, 345;
corrugata, 358; *damacornis*, 306;
episphaeria, 362; *fuliginosa*, 365;
imbricatula, 325; *Kunzei*, 323; *luteo-*
badia, 323; *rubiginosa*, 332; *setosa*,
 311; *speciosa*, 307; *tabacina*, 325

The Thelephoraceae of North America.
 IX, 177; X, 301

Thread blight, 131

Tobacco, transpiration of, 166

Tomatoes, transpiration of, 162, 163,
 164

tomentosa (*Hymenochaete*), 368

Transpiration: effect of anaesthetics on,
 234; The effect of Bordeaux mixture
 on the rate of, 153

Tuber virens, 28

U

- umbrinum* (*Stereum*), 367, 368
ungulata (*Hymenochaete*), 338, 372
unicolor (*Hymenochaete*), 342

V

- vagum* (*Corticium*), 127, 128
vagum var. *Solani* (*Corticium*), 128
vitellinum (*Stereum*), 190

W

- Weirii* (*Aleurodiscus*), 203

Z

Zeller, S. M.: Correlation of the
 strength and durability of southern
 pine, 109; and Dodge, C. W., Gau-
 tieria in North America, 133; Rhizo-
 pogon in North America, 1